



Zheng, J., Erzurumluoglu, M., Elsworth, B., Howe, L., Haycock, P., Hemani, G., Tansey, K., Laurin, C., St Pourcain, B., Warrington, N. M., Finucane, H., Price, A. L., Bulik-Sullivan, B., Anttila, V., Paternoster, L., Gaunt, T., Evans, D., & Neale, B. M. (2017). LD Hub: a centralized database and web interface to perform LD score regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic correlation analysis. *Bioinformatics*, 33(2), 272-279. <https://doi.org/10.1101/051094>, <https://doi.org/10.1093/bioinformatics/btw613>, <https://doi.org/10.1093/bioinformatics/btw613>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):

[10.1101/051094](https://doi.org/10.1101/051094)

[10.1093/bioinformatics/btw613](https://doi.org/10.1093/bioinformatics/btw613)

[10.1093/bioinformatics/btw613](https://doi.org/10.1093/bioinformatics/btw613)

[Link to publication record in Explore Bristol Research](#)

PDF-document

This is the final published version of the article (version of record). It first appeared online via Oxford University Press at <http://bioinformatics.oxfordjournals.org/content/early/2016/09/22/bioinformatics.btw613>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Text S1. References for GWAS data included in the LD Hub v1.0

- Anderson, C. et al. (2011) Meta-analysis identifies 29 additional ulcerative colitis risk loci, increasing the number of confirmed associations to 47. *Nature Genetics*, 43, 246-252.
- Bentham, J. et al. (2015) Genetic association analyses implicate aberrant regulation of innate and adaptive immunity genes in the pathogenesis of systemic lupus erythematosus. *Nature Genetics*, 47, 1457-1464.
- Benyamin, B. et al. (2013) Childhood intelligence is heritable, highly polygenic and associated with FBNP1L. *Molecular Psychiatry*, 19, 253-258.
- Berndt, S. et al. (2013) Genome-wide meta-analysis identifies 11 new loci for anthropometric traits and provides insights into genetic architecture. *Nature Genetics*, 45, 501-512.
- Boger, C. et al. (2011) CUBN Is a Gene Locus for Albuminuria. *Journal of the American Society of Nephrology*, 22, 555-570.
- Boraska, V. et al. (2014) A genome-wide association study of anorexia nervosa. *Molecular Psychiatry*, 19, 1085-1094.
- Bradfield, J. et al. (2012) A genome-wide association meta-analysis identifies new childhood obesity loci. *Nature Genetics*, 44, 526-531.
- Cordell, H. J. et al. (2015). International genome-wide meta-analysis identifies new primary biliary cirrhosis risk loci and targetable pathogenic pathways. *Nat Commun*. 6:8019.
- Dastani Z et al. (2012). Novel loci for adiponectin levels and their influence on type 2 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals. *PLoS Genet*. 8(3):e1002607.
- de Moor, M. et al. (2010) Meta-analysis of genome-wide association studies for personality. *Molecular Psychiatry*, 17, 337-349.
- Dubois, P. et al. (2010) Multiple common variants for celiac disease influencing immune gene expression. *Nature Genetics*, 42, 295-302.
- Dupuis, J. et al. (2010) New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nature Genetics*, 42, 105-116.
- Estrada, K. et al. (2012) Genome-wide meta-analysis identifies 56 bone mineral density loci and reveals 14 loci associated with risk of fracture. *Nature Genetics*, 44, 491-501.
- Frank, A. et al. (2010) Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. *Nature Genetics*, 42, 1118-1125.
- Furberg, H. et al. (2010) Genome-wide meta-analyses identify multiple loci associated with smoking behavior. *Nature Genetics*, 42, 441-447.
- Horikoshi, M. et al. (2012) New loci associated with birth weight identify genetic links between intrauterine growth and adult height and metabolism. *Nature Genetics*, 45, 76-82.
- Huffman, J. et al. (2015) Modulation of Genetic Associations with Serum Urate Levels by Body-Mass-Index in Humans. *PLOS ONE*, 10, e0119752.
- Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis (2013) *The Lancet*, 381, 1371-1379.
- KÄttgen, A. et al. (2010) New loci associated with kidney function and chronic kidney disease. *Nature Genetics*, 42, 376-384.
- Lambert, J. et al. (2013) Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nature Genetics*, 45, 1452-1458.
- Lango Allen, H. et al. (2010) Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature*, 467, 832-838.
- Liu, J. Z. et al. (2015). Association analyses identify 38 susceptibility loci for inflammatory bowel disease and highlight shared genetic risk across populations. *Nat Genet*. 47(9):979-86.
- Mahajan, A. et al. (2014) Genome-wide trans-ancestry meta-analysis provides insight into the genetic architecture of type 2 diabetes susceptibility. *Nature Genetics*, 46, 234-244.
- Manning, A. et al. (2012) A genome-wide approach accounting for body mass index identifies genetic variants influencing fasting glycemic traits and insulin resistance. *Nature Genetics*, 44, 659-669.
- Morris, A. et al. (2012) Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nature Genetics*, 44, 981-990.
- Moffatt, M. F. et al. (2007). Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature*. 448(7152):470-3.
- Neale, B. et al. (2010) Meta-Analysis of Genome-Wide Association Studies of Attention-Deficit/Hyperactivity Disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 49, 884-897.
- Okada, Y. et al. (2013) Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature*, 506, 376-381.
- Perry, J. et al. (2014) Parent-of-origin-specific allelic associations among 106 genomic loci for age at menarche. *Nature*, 514, 92-97.
- Prokopenko, I. et al. (2014) A Central Role for GRB10 in Regulation of Islet Function in Man. *PLoS Genetics*, 10, e1004235.
- Rietveld, C. et al. (2014) Common genetic variants associated with cognitive performance identified using the proxy-phenotype method. *Proceedings of the National Academy of Sciences*, 111, 13790-13794.
- Rietveld, C. et al. (2013) GWAS of 126,559 Individuals Identifies Genetic Variants Associated with Educational Attainment. *Science*, 340, 1467-1471.
- Ripke, S. et al. (2012) A mega-analysis of genome-wide association studies for major depressive disorder. *Molecular Psychiatry*, 18, 497-511.
- Ripke, S. et al. (2014) Biological insights from 108 schizophrenia-associated genetic loci. *Nature*, 511, 421-427.
- Sawcer, S. et al. (2011) Genetic risk and a primary role for cell-mediated immune mechanisms in multiple sclerosis. *Nature*, 476, 214-219.
- Saxena, R. et al. (2010) Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge. *Nature Genetics*, 42, 142-148.
- Schunkert, H. et al. (2011) Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nature Genetics*, 43, 333-338.
- Shin, S. et al. (2014) An atlas of genetic influences on human blood metabolites. *Nature Genetics*, 46, 543-550.
- Shungin, D. et al. (2015) New genetic loci link adipose and insulin biology to body fat distribution. *Nature*, 518, 187-196.
- SimÄn-SÄnchez, J. et al. (2009) Genome-wide association study reveals genetic risk underlying Parkinson's disease. *Nature Genetics*, 41, 1308-1312.
- Sklar, P. et al. (2011) Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4. *Nature Genetics*, 43, 977-983.
- Soranzo, N. et al. (2010) Common Variants at 10 Genomic Loci Influence Hemoglobin A1C Levels via Glycemic and Nonglycemic Pathways. *Diabetes*, 59, 3229-3239.
- Speliotes, E. et al. (2010) Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature Genetics*, 42, 937-948.
- Stahl, E. et al. (2010) Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. *Nature Genetics*, 42, 508-514.
- Taal, H. et al. (2012) Common variants at 12q15 and 12q24 are associated with infant head circumference. *Nature Genetics*, 44, 532-538.
- Teslovich, T. et al. (2010) Biological, clinical and population relevance of 95 loci for blood lipids. *Nature*, 466, 707-713.
- van den Berg, S. et al. (2014) Harmonization of Neuroticism and Extraversion phenotypes across inventories and cohorts in the Genetics of Personality Consortium: an application of Item Response Theory. *Behav Genet*, 44, 295-313.
- van der Valk, R. et al. (2014) A novel common variant in DCST2 is associated with length in early life and height in adulthood. *Human Molecular Genetics*, 24, 1155-1168.
- Yang, J. et al. (2012) FTO genotype is associated with phenotypic variability of body mass index. *Nature*, 490, 267-272.

